

Respiratory Viruses in Luxembourg (ReViLux)


Weekly report (27 – 03 July 2022)

Executive summary

The sentinel network reported 4.6% consultations for influenza-like illness, thus exceeding the baseline circulation threshold, according to the European Centre for Disease Prevention and Control (ECDC). Within the specimens collected by the sentinel network, SARS-CoV-2 was the most frequently detected virus over the last four weeks (51.3%), followed by Human rhinovirus (25.0%) and Parainfluenzavirus (10.5%).

Regarding the SARS-CoV-2 genomic surveillance, LNS sequenced 646 specimens from residents in Luxembourg in week 26 (of 5741 total cases in the Grand Duchy of Luxembourg; 11.3%). This exceeds the ECDC recommendations to detect emerging variants.

The Omicron variant remains the only one detected in the representative sample. The Omicron BA.5 lineage is the most frequent one (84.5%, CI 81.4 - 87.5%), followed by BA.2 (8.5%, CI 6.2 - 10.9%).



Introduction

The Laboratoire national de santé, as **National Reference Laboratory for Acute Respiratory Infections in Luxembourg**, performs close surveillance on respiratory viruses, with a special focus on SARS-CoV-2. There are currently two active projects:

- **The Sentinel Surveillance Network.** It provides a broad picture of respiratory diseases affecting the Luxembourgish population, based on its double monitoring system (syndromic and virological).
- **The National SARS-CoV-2 Genomic Surveillance Program.** It enables detailed observation of SARS-CoV-2 mutations and variants through time and space, and also monitoring specific groups of interest.

The objective of the ReViLux is to inform public health actions in Luxembourg.

Sentinel Surveillance Network

The **Sentinel Surveillance Network** aims at monitoring the circulating respiratory viruses, including SARS-CoV-2, and hence underpin public health actions. Following the World Health Organization (WHO) and European Centre for Disease Prevention and Control (ECDC) guidance, it focuses on cases of acute respiratory infection (ARI) and influenza-like illness (ILI).

Season 2021/2022 ended by week 20, and we are currently in the interseason period. The history of ILI consultations is displayed in Figure 1, and a detailed summary of the number of ARI and ILI cases during the last four weeks is included in Table 1. In the week of study, 4.6% of consultations were reported as ILI, thus exceeding the threshold for baseline circulation during the interseason, according to ECDC (1.14%).

Table 1. Syndromic surveillance over the last 4 weeks

Week	ARI		ILI		Total consultations
	N	%	N	%	
2022/23	15	8.88	7	4.14	169
2022/24	55	15.80	11	3.16	348
2022/25	54	14.17	10	2.62	381
2022/26	46	12.47	17	4.61	369

ARI: Acute Respiratory Infections; ILI: Influenza-Like Illness.




Figure 1. Percentage of patients with influenza-like illness over the last three seasons

Additionally, a selection of sentinel cases is further studied in order to monitor the circulation of respiratory viruses in the country, as shown in Figure 2. Over the last 4 weeks, the positivity rate was at 61.7%, and the most frequently detected viruses were SARS-CoV-2 (51.3%), Human rhinovirus (25.0%) and Parainfluenzavirus (10.5%). Co-infections were detected in 5 specimens, including 3 with SARS-CoV-2. These results are displayed more in detail in Table 2.




Figure 2. Distribution of respiratory viruses detected within the Sentinel Network, by two-week periods. Results from last weeks are not yet consolidated.

Co-inf: co-infection; FLU-A: influenza A; FLU-B: influenza B; PIV: parainfluenzavirus; RSV: respiratory syncytial virus; ADV: adenovirus; MPV: metapneumovirus; HRV: human rhinovirus; SC2: SARS-CoV-2.

Table 2. Distribution of respiratory viruses detected within the Sentinel Network over the last 4 weeks and during the current season.

Virus	Last 4 weeks		Interseason	
	N*	%	N*	%
SARS-CoV-2	39	51.3	40	47.1
Human rhinovirus	19	25.0	24	28.2
Parainfluenzavirus	8	10.5	9	10.6
Adenovirus	5	6.6	6	7.1
Respiratory syncytial virus	3	3.9	3	3.5
Metapneumovirus	1	1.3	2	2.4
Influenzavirus A	1	1.3	1	1.2
Influenzavirus B	0	0.0	0	0.0
Total	76	100.0	85	100.0

Co-infection cases counted once for each virus detected.

SARS-CoV-2 Genomic Surveillance

The current sequencing strategy

The National Reference Laboratory for Acute Respiratory Infections at LNS receives SARS-CoV-2 positive samples (nasopharyngeal or oropharyngeal swabs analysed by RT-PCR) from the national network of laboratories and proceeds as follows:

1. Sequencing a representative sample of specimens.
2. Sequencing specimens from target groups (i.e. hospital cases and post-vaccination cases).
3. Sequencing specimens from clusters with high transmission.

The representative sample of specimens is a systematic selection from all SARS-CoV-2 positive cases registered in Luxembourg to detect emerging variants and early increases in their incidence and transmission within the community in Luxembourg. This sample is selected according to the ECDC guidelines.

A screening by targeted PCR tests is also carried out when a new variant emerges in order to enable earlier evaluation of its spread in the population.

SARS-CoV-2 lineages have been assigned based on Rambaut et al. using the Phylogenetic Assignment of Named Global Outbreak LINeages (pangolin) software (4.1.1, designation v1.11). The Pango nomenclature is used in addition to the WHO nomenclature to enable easier visualization of links between any evolving variants and their ancestor.

Sequencing activity



Figure 3. Flowchart of specimens collected during week 26/2022

In week 26, 5741 new cases were registered in Luxembourg; hence, the minimum sample size required to detect emerging variants at a 2.5% incidence is estimated to be 543 specimens (9.5%).

As shown in Figure 3, the microbial genomics unit at the LNS sequenced 728 specimens from the week of study, including 646 national ones. The weekly sequencing coverage remains at 11.3% (out of 5741 cases registered in Luxembourg; see coverage trend in Figure 4), which exceeds the recommended sample size.




Figure 4. Number of specimens sequenced and national coverage based on weekly number of positive cases in Luxembourg. The coverage from the last two weeks is not consolidated yet.

Circulating lineages detection

The distribution of successfully assigned lineages within the national selection is shown in Figure 6, and it is further detailed in Table 4 (last two weeks). A summary of the VOCs assigned among all specimens sequenced (including non-residents) during the last two weeks and since the beginning of the sequencing activity is shown in Table 5.

The Omicron variant remains the only one within the representative sample, the most frequent lineage being Omicron BA.5 (84.5%), followed by followed by BA.2 (8.5%).

Table 4. Distribution of SARS-CoV-2 lineages detected within the representative sample during the last two weeks. Previously reported cases might be updated by retrospective analysis.

Lineage	Previous week			Current week		
	N	%	CI %	N	%	CI %
Omicron BA.5	409	77.3	73.7 - 80.9	457	84.5	81.4 - 87.5
Omicron BA.2	84	15.9	12.8 - 19.0	46	8.5	6.2 - 10.9
Omicron BA.4	35	6.6	4.5 - 8.7	38	7.0	4.9 - 9.2
Omicron BA.1	1	0.2	0.0 - 0.6	0	0.0	-
Total	529	100.0	-	541	100.0	-

CI: Confidence Interval at 95%.




Figure 6. Distribution of lineages within the representative sample during the last 10 weeks.

Table 5. Distribution SARS-CoV-2 variants of concern and lineages under monitoring detected within all samples sequenced since the beginning of the pandemic.

Variant	Previous week		Current week		Cumulative count
	N	%	N	%	
Omicron	576	100.0	600	100.0	19 468
BA.5	244	42.4	218	36.3	1 393
BA.5.1	36	6.3	88	14.7	239
BA.2.12.1	15	2.6	10	1.7	214
BA.4	28	4.9	25	4.2	155
BA.2.13	0	0.0	0	0.0	43
BA.2.11	0	0.0	0	0.0	2
BA.2.75	0	0.0	0	0.0	0
Delta	0	0.0	0	0.0	13 679
Others	0	0.0	0	0.0	15 005
Total	576	100.0	600	100.0	48 152

Clinical and epidemiological factors

In this section, the lineage distribution of all specimens sequenced over the last month is assessed by demographics (sex and age group, Figure 7) and sampling setting (community vs. hospital, Table 6).




Figure 7. Age group and sex distribution of specimens sequenced over the last 4 weeks, by lineage.

Table 6. Comparison of lineage distribution by sampling setting.

Lineage	Community			Hospital		
	Women	Men	Total	Women	Men	Total
Omicron BA.5	68.9%	75.6%	71.9%	82.5%	82.9%	82.7%
Omicron BA.2	21.2%	18.5%	20.0%	13.3%	12.8%	13.1%
Omicron BA.4	9.9%	5.9%	8.1%	4.2%	4.3%	4.2%
Total	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%

Mutation surveillance

In addition to the surveillance of SARS-CoV-2 variants, the LNS monitors the occurrence of SARS-CoV-2 mutations reported to have a clinical and epidemiological relevance. Newly acquired mutations may occur and their early detection might be key to expect changes in the epidemic evolution. Following ECDC guidance, the LNS is currently monitoring a selection of mutations to the spike protein frequently associated to VOCs. As each VOC is characterised by a set of defining mutations, which are expected to be highly present, it is interesting to analyse the non-defining ones.

Among the specimens collected over the last four weeks that could not be assigned to a known lineage, the F486V mutation (S gene) was detected in a higher proportion than expected for Europe (98.86% vs 71.72%, according to GISAID). This mutation is most frequently found in BA.5 lineages and sublineages.

References

Centers for Disease Control and Prevention. SARS-CoV-2 Variant Classifications and Definitions. Retrieved 11 July 2022, from <https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html>

COVID-19 Data Portal - accelerating scientific research through data. (2021). Retrieved 11 July 2022, from <https://www.covid19dataportal.org/sequences>

Direção-Geral da Saúde. Relatório de Situação. Retrieved 11 July 2022, from <https://covid19.min-saude.pt/>

European Centre for Disease Prevention and Control. Guidance for representative and targeted genomic SARS-CoV-2 monitoring – 3 May 2021. ECDC : Stockholm ; 2021

European Centre for Disease Prevention and Control. SARS-CoV-2 variants of concern. Retrieved 11 July 2022, from <https://www.ecdc.europa.eu/en/covid-19/variants-concern>

Genomic sequencing of SARS-CoV-2: a guide to implementation for maximum impact on public health. Geneva: World Health Organization; 2021.

GISAID. EpiCoV – Pandemic coronavirus causing COVID-19. Retrieved 5 mai 2022, from <https://www.gisaid.org>

GitHub - cov-lineages/pangolin: Software package for assigning SARS-CoV-2 genome sequences to global lineages. (2021). Retrieved 11 July 2022, from <https://github.com/cov-lineages/pangolin>

Hadfield J., Megill C., Bell S., Huddleston J., Potter B., Callender C. et al. (2018). Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics*, 34(23), 4121-4123. doi: 10.1093/bioinformatics/bty407

Instituto Nacional de Saúde Doutor Ricardo Jorge. Diversidade genética do novo coronavírus SARS-CoV-2 (COVID-19) em Portugal. Retrieved 11 July 2022, from <https://insaflu.insa.pt/covid19/>

Rambaut A., Holmes E., O’Toole Á., Hill V., McCrone J., Ruis C. et al. (2020). A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. *Nature Microbiology*, 5(11), 1403-1407. doi: 10.1038/s41564-020-0770-5

Robert Koch Institut. Aktueller Lage-/Situationsbericht des RKI zu COVID-19. Retrieved 11 July 2022, from https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Situationsberichte/Gesamt.html;jsessionid=69BC18053F9591C56EB148E463103DB7.internet101

Santé publique France. Coronavirus : chiffres clés et évolution de la COVID-19 en France et dans le Monde. Retrieved 11 July 2022, from <https://www.santepubliquefrance.fr/dossiers/coronavirus-covid-19/coronavirus-chiffres-cles-et-evolution-de-la-covid-19-en-france-et-dans-le-monde>

Sciensano. COVID-19 – Bulletin épidémiologique hebdomadaire. Retrieved 11 July 2022, from <https://covid-19.sciensano.be/fr/covid-19-situation-epidemiologique>

The Luxembourgish data platform. COVID-19: Rapports hebdomadaires. Retrieved 11 July 2022, from <https://data.public.lu/en/datasets/covid-19-rapports-hebdomadaires/>