

Respiratory Viruses in Luxembourg (ReViLux)

Weekly report (21 December 2022)

Executive summary

The sentinel network detected a very high epidemic activity over the last week, based on 14.6% of consultations being associated with influenza-like illness.

Within the specimens collected by the sentinel network, Influenzavirus A was the most frequently detected virus over the last two weeks (36.6% positivity rate), followed by Human rhinovirus (15.1%) and Respiratory syncytial virus (RSV) (13.4%). Genotyping analyses showed that the most frequent RSV strain during the current season is RSV-B (70.9%), and the most frequent influenza A is H3 (94%).

Regarding the SARS-CoV-2 genomic surveillance, The Omicron BQ.1 lineage is the most frequent one (58.8%, confidence interval: 52.7 - 65.0%), followed by Omicron BA.5 (28%, confidence interval: 22.3 - 33.6%).

IN THE SPOTLIGHT

Respiratory syncytial virus is a common cause of lower respiratory infections (bronchiolitis and pneumonia), especially in children and older adults. The first exposure to the virus and having comorbidities are risk factors for presenting more severe symptoms.

BQ.1 and its sublineages are a subtype of SARS-CoV-2 Omicron BA.5. There are no data yet suggesting higher severity than other Omicron lineages, but it carries some mutations that have been associated to escape immunity from previous BA.5 infections, which suggests a higher reinfection risk. Vaccine protection against infection might be reduced, but protection against severe disease is not expected to be affected.

XBB is a recombinant of two lineages, BA.2.10.1 and BA.2.75. Early evidence suggests higher reinfection risk compared to other circulating lineages, due to the mutations it carries in the Receptor-Binding Domain of the spike protein. Nevertheless, there is no evidence suggesting an increase in severity.

Introduction

The Laboratoire national de santé, as **National Reference Laboratory for Acute Respiratory Infections** in Luxembourg, performs close surveillance on respiratory viruses, with a special focus on SARS-CoV-2. The main monitoring activities are:

- **The Sentinel Surveillance.** It provides a broad picture of respiratory diseases affecting the Luxembourgish population, based on clinical and laboratory data.
- **The SARS-CoV-2 Genomic Surveillance.** It enables detailed observation of SARS-CoV-2 variants and mutations through time and space, as well as studying specific groups of interest.

The objective of the ReViLux report is to inform public health actions in Luxembourg.

Sentinel Surveillance Network

The Sentinel Surveillance aims at monitoring the circulating respiratory viruses, from traditional ones like Influenza to more recent ones like SARS-CoV-2, and hence underpin public health actions. The Sentinel Network is a group of general practitioners and paediatricians spread over the country. They report the weekly number of patients showing symptoms suggestive of acute respiratory infection (ARI) and influenza-like illness (ILI), and those patients are then sampled and tested for a panel of respiratory viruses. The circulation of respiratory viruses in the north hemisphere is generally monitored by seasons that go from week 40 to week 20. The period between weeks 20 and 40 is usually called interseason.

Clinical results

Last week, 14.6% of consultations were reported as ILI, which represents a very high epidemic activity for Luxembourg, according to ECDC and the Moving Epidemic Method. The history of ILI consultations is displayed in Figure 1, and a detailed summary of the number of ARI and ILI cases during the last four weeks is included in Table 1.

Laboratory results

Over the last two weeks, the most frequently detected viruses (and their positivity rates) were Influenzavirus A (36.6%), Human rhinovirus (15.1%) and Respiratory syncytial virus (13.4%). An overview of the viral circulation during the current season is displayed in Figure 2 and Table 2.




Figure 1. Percentage of patients with influenza-like illness over the last three seasons
Background colours according to intensity of circulation: low, medium, high, very high.

Table 1. Syndromic surveillance over the last 4 weeks

Week	ARI		ILI		Total consultations
	N	%	N	%	
2022/47	106	18.63	18	3.16	569
2022/48	129	21.61	36	6.03	597
2022/49	98	19.60	21	4.20	500
2022/50	82	17.15	70	14.64	478

ARI: Acute Respiratory Infections; ILI: Influenza-Like Illness.




Figure 2. Distribution of respiratory viruses detected within the Sentinel Network, by two-week periods. Results from last weeks are not yet consolidated.

FLU-A: influenza A; FLU-B: influenza B; PIV: parainfluenza virus; RSV: respiratory syncytial virus; ADV: adenovirus; MPV: metapneumovirus; HRV: human rhinovirus; SC2: SARS-CoV-2.

Table 2. Distribution of respiratory viruses detected within the Sentinel Network over the last 2 weeks compared to the interseason period.

Virus	Last 2 weeks		Season 2022/2023	
	N*	Positivity rate (%)	N*	Positivity rate (%)
Influenzavirus A	135	36.6	180	19.5
Human rhinovirus	36	15.1	163	22.1
Respiratory syncytial virus	32	13.4	145	19.7
SARS-CoV-2	39	10.6	126	15.4
Metapneumovirus	25	10.5	42	5.7
Adenovirus	22	9.2	83	11.3
Influenzavirus B	3	0.8	4	0.4
Parainfluenzavirus	1	0.4	18	2.4
Total	293		761	

*Co-infection cases counted once for each virus detected.

Genotyping analyses

In order to assess the circulation of viral strains, genotyping analyses were also performed for Influenzavirus A and Respiratory syncytial virus (RSV) (Figure 3). During the current season, the most frequently detected strain for Influenzavirus A was H3 (94%), and for RSV it was type B (70.9%).




Figure 3. Detection of influenzavirus-A (Flu-A) and Respiratory syncytial virus (RSV) strains in sentinel samples during the current season.

SARS-CoV-2 Genomic Surveillance

SARS-CoV-2 has posed a serious threat to the world population since 2020, and it is still a challenge for health systems. For this reason, the National Reference Laboratory for Acute Respiratory Infections at LNS continues to monitor the genomic characteristics of the SARS-CoV-2 virus detected in Luxembourg.

LNS receives positive specimens (nasopharyngeal or oropharyngeal swabs analysed by RT-PCR) from the national network of laboratories. All hospital specimens are sequenced, as well as a representative selection of community specimens. Illumina and Nanopore devices are used respectively. When needed, an additional screening by targeted PCR is also performed to enable earlier evaluation of the spread of newly emerging variants.

Bioinformatic analyses are based on a standardised pipeline, and lineage assignment is performed through the PANGOLin software (4.1.3, pango-data 1.16, mode USHER). In order to allow easier visualisation, WHO and Nextstrain categorisations are also used.

Sequencing activity

In week 49, 1440 new cases were registered in Luxembourg; hence, the minimum sample size required to detect emerging variants at a 5% incidence is estimated to be 243 specimens (16.9%). The microbial genomics unit at LNS sequenced 374 specimens from the week of study, including 333 national ones (sequencing coverage: 23.1%; see Figure 4). This reaches the recommended sample size.




Figure 4. Number of specimens sequenced and national coverage based on weekly number of positive cases in Luxembourg. The coverage from the last two weeks is not consolidated yet.

Variant circulation

The Omicron variant remains the main one detected within the representative sample, the most frequent subvariant being Omicron BQ.1 (58.8%), followed by Omicron BA.5 (28%).

An overview of the variants and lineages circulating over the last weeks is displayed in Figure 5, and further details are shown in Table 3. The history of the circulation of each variant since January 2021 is displayed in Figure 6.




Figure 5. Distribution of lineages within the representative sample during the last 10 weeks.

Table 3. Distribution of SARS-CoV-2 lineages detected within the representative sample during the last two weeks. Previously reported cases might be updated by retrospective analysis.

Lineage*	Week 48		Week 49	
	%	CI %	%	CI %
Omicron BQ.1	55.4	48.9 - 61.9	58.8	52.7 - 65.0
Omicron BA.5	31.1	25.0 - 37.2	28.0	22.3 - 33.6
Omicron BA.2.75	8.6	4.9 - 12.2	7.8	4.4 - 11.2
Recombinant XBB	2.3	0.3 - 4.2	4.1	1.6 - 6.6
Omicron BA.4	2.7	0.6 - 4.8	1.2	0.0 - 2.6

CI: Confidence Interval at 95%. *Sub-lineages BA.2.75 and BQ.1 are not included in the count of their parental lineages (BA.2 and BA.5, respectively).




Figure 6. Proportion of each variant circulating in Luxembourg since January 2021.

Clinical and epidemiological factors

In this section, the lineage distribution of all specimens sequenced over the last month is assessed by demographics (sex and age group, Figure 7) and sampling setting (community vs. hospital, Table 4).




Figure 7. Age group and sex distribution of specimens sequenced over the last 4 weeks, by lineage.

Table 4. Comparison of lineage distribution by sampling setting.

Lineage	Community			Hospital		
	Women	Men	Total	Women	Men	Total
Omicron BQ.1	65.2%	64.1%	64.7%	58.5%	61.4%	59.8%
Omicron BA.5	34.8%	35.9%	35.3%	41.5%	38.6%	40.2%
Total	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%

Mutation surveillance

In addition to the surveillance of SARS-CoV-2 variants, the LNS monitors the occurrence of SARS-CoV-2 mutations, as their early detection might be key to foresee changes in the epidemic evolution. The LNS is currently monitoring mutations to the spike protein, following ECDC guidance, and comparing their prevalence to that observed in Europe (according to GISAID).

Among the specimens collected over the last four weeks, we detected a higher circulation of the following mutations in Luxembourg (compared to Europe):

Table 5. Mutations detected at a higher proportion in Luxembourg

Lineage	Mutation	Gene	% in LU	% in EU	Ratio
BA.5	K444T	S	63.55	56.5	1.13

LU: Luxembourg; EU: Europe

References

Centers for Disease Control and Prevention. SARS-CoV-2 Variant Classifications and Definitions. Retrieved 21 December 2022, from <https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html>

European Centre for Disease Prevention and Control. Guidance for representative and targeted genomic SARS-CoV-2 monitoring – 3 May 2021. ECDC : Stockholm ; 2021

European Centre for Disease Prevention and Control. SARS-CoV-2 variants of concern. Retrieved 21 December 2022, from <https://www.ecdc.europa.eu/en/covid-19/variants-concern>

Genomic sequencing of SARS-CoV-2: a guide to implementation for maximum impact on public health. Geneva: World Health Organization; 2021.

GISAID. EpiCoV – Pandemic coronavirus causing COVID-19. Retrieved 5 mai 2022, from <https://www.gisaid.org>

GitHub - cov-lineages/pangolin: Software package for assigning SARS-CoV-2 genome sequences to global lineages. (2021). Retrieved 21 December 2022, from <https://github.com/cov-lineages/pangolin>

Hadfield J., Megill C., Bell S., Huddleston J., Potter B., Callender C. et al. (2018). Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics*, 34(23), 4121-4123. doi: 10.1093/bioinformatics/bty407

Rambaut A., Holmes E., O’Toole Á., Hill V., McCrone J., Ruis C. et al. (2020). A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. *Nature Microbiology*, 5(11), 1403-1407. doi: 10.1038/s41564-020-0770-5

The Luxembourgish data platform. COVID-19: Rapports hebdomadaires. Retrieved 21 December 2022, from <https://data.public.lu/en/datasets/covid-19-rapports-hebdomadaires/>