

# Respiratory Viruses in Luxembourg (ReViLux)

## Sentinel Network Report -Week 11

### Summary of Sentinel Network activities

At the end of week **2025/11**, the sentinel network detected a medium epidemic activity, based on **11.3%** of consultations being associated with influenza-like illness. Among the specimens collected by the sentinel network over the last week, the percentage of positive tests for **Human rhinovirus** was **15.1%**, **13.1%** for **Influenza B** and **10.3%** for **Influenza A**, while **RSV** activity was at **1.9%** and that of **SARS-CoV-2** at a very low level.

In total, this season (24/25) 2,491 samples were received with 855 Influenza positive samples (381 Influenza B and 474 Influenza A). So far, 452 of the 474 Influenza A samples (95.4%) have been subtyped. Hundred and ninety four (42.9%) Influenza A samples have been subtyped as A(H1)pdm09 and 258 (57.1%) samples as A(H3) virus. Combined Influenza A and B positivity rates decreased from 30.8% (2025/10) to 23.4% (2025/11), but still indicate a mixture of subtypes spread across the country.

RSV continues to circulate at low level (1.9%) with 41.8% of subtype A and 58.2% subtype B cases.

Syndromic surveillance over the last 4 weeks (Table 1)

Week	ARI		ILI		Total consultations
	N	%	N	%	
2025/08	70	19.94	48	13.68	351
2025/09	42	10.97	38	9.92	383
2025/10	60	16.71	34	9.47	359
2025/11	36	11.32	36	11.32	318

*ARI: Acute Respiratory Infections; ILI: Influenza-Like Illness.*

---

## Sentinel Surveillance Network

The Sentinel Surveillance aims to monitor circulating respiratory viruses, from traditional ones like Influenza to more recent ones like SARS-CoV-2, and hence underpin public health actions. The Sentinel Network is a group of general practitioners and paediatricians spread across the country. They report the weekly number of patients showing symptoms suggestive of acute respiratory infection (ARI) and influenza-like illness (ILI), and those patients are then sampled and tested for a panel of respiratory viruses. The circulation of respiratory viruses in the north hemisphere is generally monitored by seasons that go from week 40 to week 20. The period between weeks 20 and 40 is usually called inter-season.

### Clinical results

Last week (**end of week 2025/11**), **11.3%** of the consultations were reported as ILI, representing a medium epidemic activity for Luxembourg, according to ECDC and the Moving Epidemic Method. Over the last four weeks medium ILI rates have been observed in Luxembourg. The ILI rate increased slightly from 9.5% (2025/10) to 11.3% (2025/11), still indicating considerable respiratory virus activity.

The history of ILI consultations is displayed in figure 1, and a detailed summary of the number of ARI and ILI cases during the last four weeks is included in table 1.

### Laboratory results

Over the last week, the most frequently detected viruses (according to positivity rates) were **Human rhinovirus (15.1%)**, followed by **Influenza B (13.1%)** and **Influenza A (10.3%)**. During week 2025/11, Influenza B decreased slightly to 13%, Influenza A decreased from 16.3% (2025/10) to 10.3%, while SARS-CoV-2 activity remained at very low level within the sentinel network. Overall Influenza positivity decreased from around 30% (2025/10) to 23.4% (2025/11) (table 2 and figure 4). An overview of the circulating viral pathogens during the current and previous inter- season is displayed in figure 2 and table 2.

More than 70% of the Influenza B cases identified this week were in children under 5 years of age, while Influenza A has been detected in all age-groups under 65 years old and all cases have been subtyped as A(H3) virus so far.

Over the last week (2025/11), RSV activity decreased to 2% and remained below 5% for two consecutive weeks. So far this season (24/25), two hundred and sixty eight RSV cases have been detected, including ninety seven RSV-A cases and hundred and thirty five RSV-B cases. Approximately, 55% of cases were aged 1 and 4 years (figure 3).

In total, 2491 sentinel samples have been analysed with more than 60% of samples belonging to age-group below 18 years and with 52% of female cases. Over the past week, 35% of samples were from adults over 18 years old and 4% from adults above 65 years old (figure 4).

Over the past 2 weeks, Human rhinovirus was detected in all age-groups, while Adenovirus and Metapneumovirus were detected predominantly in children. Over the last 2 weeks, about 84% of all co-infections (N=19) were detected primarily in children below 5 years and 16% in children aged between 5 and 18 years. The most commonly identified combination was Human rhinovirus with Metapneumovirus. So far this season, ten patients were co-infected with Influenza A and B.




Figure 1. Percentage of patients with Influenza-like illness over the last three seasons  
Background colours according to intensity of circulation: baseline, low, medium, high, very high.

Table 2. Distribution of respiratory viruses detected within the Sentinel Network previous 4 weeks compared to previous year. Data for week 2025/11 not yet completely consolidated

Virus	Season 2024/25					Season 2023/24	
	Positivity Rate in %					Positivity Rate in %	
	W08	W09	W10	W11	Total N (%)	W11	Total N (%)
Human rhinovirus	17.0	15.9	25.0	15.1	580 (23.4)	37.3	572 (24.9)
Influenzavirus B	33.3	19.6	14.4	13.1	381 (15.3)	0.0	12 (0.5)
Influenzavirus A	12.6	19.6	16.3	10.3	474 (19.0)	4.8	388 (16.5)
Adenovirus	7.4	4.7	9.6	10.4	159 (6.4)	6.8	125 (5.4)
Metapneumovirus	3.0	6.5	7.7	9.4	90 (3.8)	25.4	125 (5.4)
Parainfluenzavirus	0.7	0.0	1.0	2.8	66 (2.7)	3.4	77 (3.4)
Respiratory syncytial virus	7.4	5.6	4.8	1.9	268 (10.8)	0.0	212 (9.2)
SARS-CoV-2	1.5	0.9	0.0	0.9	70 (2.8)	1.6	227 (9.7)

Co-detection counted once for each virus detected & N- total number of detection during season




Figure 2. Distribution of respiratory viruses detected within the Sentinel Network, by calendar week. Results from last weeks are not all yet consolidated.  
 FLU-A: Influenzavirus A; FLU-B: Influenzavirus B; PIV: Parainfluenzavirus; RSV: Respiratory syncytial virus; ADV: Adenovirus; MPV: Metapneumovirus; HRV: Human rhinovirus; SC2: SARS-CoV-2.

Figure 3. Displays RSV cases according to different age groups with highest impact among the 1 years old. Data for week 2025/11 not yet completely consolidated




Figure 4. Displays number of sentinel samples received per week by age-group including overall sample positivity including Human rhinovirus (HRV, dotted line), excluding HRV (black line) and Influenza combined (red). Secondary axis corresponds to positivity





Figure 5. Displays detection of Influenza subtypes by age-group. Data for week 2025/11 not yet completely consolidated



## References

European Centre for Disease Prevention and Control. SARS-CoV-2 variants of concern. Retrieved 18 March 2025. from <https://www.ecdc.europa.eu/en/covid-19/variants-concern>

European Centre for Disease Prevention and Control. Communicable Disease Threats Report Week 8-14-March-2025-week-11 <https://www.ecdc.europa.eu/en/publications-data/communicable-disease-threats-report-8-14-march-2025-week-11>

GISAID. EpiCoV – Pandemic coronavirus causing COVID-19. Retrieved 18 March 2025. from <https://www.gisaid.org>

GitHub - cov-lineages/pangolin: pango-designation/lineage\_notes.txt. (2025). Retrieved 04 March 2025. from [https://github.com/cov-lineages/pango-designation/blob/master/lineage\\_notes.txt](https://github.com/cov-lineages/pango-designation/blob/master/lineage_notes.txt)

Hadfield J. Megill C. Bell S. Huddleston J. Potter B. Callender C. et al. (2018). Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics*. 34(23). 4121-4123. doi: 10.1093/bioinformatics/bty407

Rambaut A. Holmes E. O’Toole Á. Hill V. McCrone J. Ruis C. et al. (2020). A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. *Nature Microbiology*. 5(11). 1403-1407. doi: 10.1038/s41564-020-0770-5

World Health Organization. Tracking sars-COV-2 variants Retrieved 04 March 2025. from <https://www.who.int/activities/tracking-SARS-CoV-2-variants>

World Health Organization. Risk evaluation for SARS-CoV-2 Variant Under Monitoring: LP.8.1. from <https://www.who.int/publications/m/item/risk-evaluation-for-sars-cov-2-variant-under-monitoring-lp81>